arXiv:1205.6547v1 [math.NT] 30 May 2012

HERMITE POLYNOMIALS RELATED TO GENOCCHI, EULER AND BERNSTEIN POLYNOMIALS

SERKAN ARACI, **JONG JIN SEO**, AND **MEHMET ACIKGOZ**

Abstract. The objective of this paper is to derive some interesting properties of Genocchi, Euler and Bernstein polynomials by means of the orthogonality of Hermite polynomials.

2010 Mathematics Subject Classification. 11S80, 11B68.

Keywords and phrases. Genocchi numbers and polynomials, Euler numbers and polynomials, Bernstein polynomials, orhogonality.

1. **Introduction**

The Genocchi, Euler and Bernstein polynomials possess many interesting properties and arising in many areas of mathematics. These polynomials have been studied by many researcher (see [1-17]).

Recently, Kim *et al.*, studied on the Hermite polynomials and their applications associated with Bernoulli an Euler numbers in "D. S. Kim, T. Kim, S. H. Rim and S. H. Lee., Hermite polynomials and their applications associated with Bernoulli and Euler numbers, Discrete Dynamics in Nature and Society, http://www.hindawi.com/journals/ddns/aip/974632/ (Article in Press). They derived some interesting properties of Hermite polynomials by using its orthogonality properties. They also showed that Hermite polynomials related to Bernoulli and Euler polynomials. It is objective of this paper to derive for Bernstein, Genocchi and Euler polynomials by using same method of theirs.

We firstly list definition of Euler, Genocchi, Bernstein and Hermite polynomials as follows:

The ordinary Euler polynomials are defined by the means of the following generating function:

$$\sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!} = e^{xt}\frac{2}{e^t+1}. \tag{1}$$

Substituting $x=0$ into (1), then we have $E_n(0) := E_n$, which is called the Euler numbers (for details, see [1], [3], [5], [10], [11] and [12]).

As is well-known, Genocchi polynomials are also defined by

$$\sum_{n=0}^{\infty} G_n(x)\frac{t^n}{n!} = e^{xt}\frac{2t}{e^t+1}. \tag{2}$$

Similarly, for $x=0$ in (2), $G_n(0) := G_n$ are Genocchi numbers (for details about this subject, see [12], [16] and [17]).

Let $C\left([0,1]\right)$ be the space of continuous functions on $[0,1]$. For $C\left([0,1]\right)$, the Bernstein operator for f is defined by

$$B_n\left(f,x\right)=\sum_{k=0}^{n}f\left(\frac{k}{n}\right)B_{k,n}\left(x\right)=\sum_{k=0}^{n}f\left(\frac{k}{n}\right)\binom{n}{k}x^k\left(1-x\right)^{n-k}$$

where $n,\ k\in\mathbb{Z}_+:=\{0,1,2,3,...\}$. Here $B_{k,n}\left(x\right)$ is called Bernstein polynomials, which are defined by

$$B_{k,n}\left(x\right)=\binom{n}{k}x^k\left(1-x\right)^{n-k},\ x\in[0,1] \tag{3}$$

(see [1], [4] and [17]).

The Hermite polynomials are defined by the generating function as follows:

$$\sum_{n=0}^{\infty}H_n\left(x\right)\frac{t^n}{n!}=e^{H(x)t}=e^{2tx-t^2} \tag{4}$$

with the usual convention about replacing by $H^n\left(x\right)$ by $H_n\left(x\right)$.

The Hermite polynomials have the following properties

$$H_n\left(x\right)=\left(-1\right)^n e^{x^2}\frac{d^n e^{-x^2}}{dx^n}. \tag{5}$$

By (4), we have the following

$$\frac{dH_n\left(x\right)}{dx}=2nH_{n-1}\left(x\right),\ n\in\mathbb{N}. \tag{6}$$

The Hermite polynomials have the orthogonal properties as follows:

$$\int_{-\infty}^{\infty}e^{-x^2}H_n\left(x\right)H_m\left(x\right)dx=\begin{cases}2^n n!\sqrt{\pi}, \text{ if } n=m\\ 0, \text{ if } n\neq m.\end{cases} \tag{7}$$

By (7), it is not difficult to show that

$$\int_{-\infty}^{\infty}e^{-x^2}x^l dx=\begin{cases}0, \text{ if } l\equiv 1\,(\mathrm{mod}\,2)\\ \frac{l!\sqrt{\pi}}{2^l\left(\frac{l}{2}\right)!}, \text{ if } l\equiv 0(\mathrm{mod}\,2),\end{cases} \tag{8}$$

where $l\in\mathbb{Z}_+$. By (8), we can derive the following

$$\int_{-\infty}^{\infty}\left(\frac{d^n e^{-x^2}}{dx^n}\right)x^m dx=\begin{cases}0, \text{ if } n>m \text{ with } n-m\equiv 1(\mathrm{mod}\,2)\\ \frac{m!(-1)^n\sqrt{\pi}}{2^{m-n}\left(\frac{m-n}{2}\right)!}, \text{ if } n\leq m \text{ with } n-m\equiv 0(\mathrm{mod}\,2).\end{cases} \tag{9}$$

We note that $H_o\left(x\right),H_1\left(x\right),H_2\left(x\right),...,H_n\left(x\right)$ are orthogonal basis for the space $\mathcal{P}_n=\{p\left(x\right)\in\mathbb{Q}\left[x\right]\mid \deg p\left(x\right)\leq n\}$ with respect to the inner product

$$\left\langle p\left(x\right),q\left(x\right)\right\rangle=\int_{-\infty}^{\infty}e^{-x^2}p(x)q\left(x\right)dx. \tag{10}$$

For $p\left(x\right)\in\mathcal{P}_n$, the polynomial $p\left(x\right)$ is given by

$$p(x)=\sum_{k=0}^{n}C_kH_k\left(x\right) \tag{11}$$

where

$$C_k = \frac{1}{2^k k!\sqrt{\pi}}\langle p(x), H_k(x)\rangle = \frac{(-1)^k}{2^k k!\sqrt{\pi}}\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)p(x)\,dx. \tag{12}$$

For more informations of Eqs. (4-12), you can refer to [11].

2. Main Results

In this section, we consider families of Bernoulli, Bernstein and Genocchi polynomials. Then, we discover novel properties for them.

Let us take $p(x) = G_n(x)$. From (11), $p(x)$ can be rewritten as

$$G_n(x) = \sum_{k=0}^{n} C_k H_k(x)$$

where

$$C_k = \frac{(-1)^k}{2^k k!\sqrt{\pi}}\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)G_n(x)\,dx.$$

Now, let us solve $\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)G_n(x)\,dx$ as follows:

$$\begin{aligned}\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)G_n(x)\,dx &= (-n)(-(n-1))\dots(-(n-k+1))\int_{-\infty}^{\infty}e^{-x^2}G_{n-k}(x)\,dx\\ &= \frac{n!}{2^k k!}\sum_{\substack{0\le l\le n-k\\ l\equiv 0 \pmod 2}}\frac{G_{n-k-l}}{(n-k-l)!2^l\left(\frac{l}{2}\right)!}.\end{aligned}$$

Thus, we procure the following theorem.

Theorem 2.1. *For $n \in \mathbb{Z}_+$, we have*

$$G_n(x) = n!\sum_{k=0}^{n}\frac{1}{2^k k!}\sum_{\substack{0\le l\le n-k\\ l\equiv 0 \pmod 2}}\frac{G_{n-k-l}H_k(x)}{(n-k-l)!2^l\left(\frac{l}{2}\right)!}.$$

Now we consider $p(x) = B_{l,n}(x)$. From (11), we note that, $p(x)$ can be rewritten as

$$B_{l,n}(x) = \sum_{k=0}^{n} C_k H_k(x)$$

where

$$\begin{aligned}C_k &= \frac{(-1)^k}{2^k k!\sqrt{\pi}}\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)B_{l,n}(x)\,dx\\ &= \frac{(-1)^k}{2^k k!\sqrt{\pi}}\frac{n!}{l!(n-l)!}\left[\sum_{j=0}^{n-l}\frac{(n-l)!}{j!(n-l-j)!}(-1)^j\left\{\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)x^{l+j}dx\right\}\right]\\ &= \frac{n!}{k!}\sum_{\substack{0\le j\le n-l\\ k-l-j\equiv 0 \pmod 2}}\binom{l+j}{l}\frac{(-1)^j}{(n-l-j)!2^{l+j}\left(\frac{l+j-k}{2}\right)!}.\end{aligned}$$

Thus, we have novel properties of Bernstein polynomials with the following theorem.

Theorem 2.2. *For $n \in \mathbb{Z}_+$ and $0 \leq l \leq n$, we have*

$$B_{l,n}(x) = n!\sum_{k=0}^{n}\frac{1}{k!}\sum_{\substack{0\leq j\leq n-l \\ k-l-j\equiv 0 (\operatorname{mod} 2)}}\binom{l+j}{l}\frac{(-1)^{j}}{(n-l-j)!2^{l+j}\left(\frac{l+j-k}{2}\right)!}H_k(x).$$

In [10], Kim *et al.*, derived the following interesting equality:

$$\sum_{k=0}^{n}E_k(x)x^{n-k} = \frac{1}{2}\sum_{j=0}^{n-1}\binom{n+1}{j}\left\{-\sum_{l=j}^{n-1}E_{l-j}+2\right\}E_j(x)+(n+1)E_n(x).$$

We now consider as $p(x) = \sum_{k=0}^{n}E_k(x)x^{n-k}$. Then, we compute as follows:

$$\begin{aligned}
C_k &= \frac{(-1)^k}{2^k k!\sqrt{\pi}}\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)p(x)\,dx \\
&= \frac{(-1)^k}{2^k k!\sqrt{\pi}}\left\{\begin{array}{c} \frac{1}{2}\sum_{j=0}^{n-1}\binom{n+1}{j}\left\{-\sum_{l=j}^{n-1}E_{l-j}+2\right\}\left[\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)E_j(x)\,dx\right] \\ +(n+1)\int_{-\infty}^{\infty}\left(\frac{d^k e^{-x^2}}{dx^k}\right)E_n(x)\,dx \end{array}\right\} \\
&= \frac{(-1)^k}{2^k k!\sqrt{\pi}}\left\{\begin{array}{c} \frac{1}{2}\sum_{j=0}^{n-1}\binom{n+1}{j}\left\{-\sum_{l=j}^{n-1}E_{l-j}+2\right\}\left[\frac{(-1)^k j!}{(j-k)!}\int_{-\infty}^{\infty}e^{-x^2}E_{j-k}(x)\,dx\right] \\ +(n+1)\frac{(-1)^n n!}{(n-k)!}\int_{-\infty}^{\infty}e^{-x^2}E_{n-k}(x)\,dx \end{array}\right\}
\end{aligned}$$

After these applications, we easily reach the following

$$\begin{aligned}
C_k &= \frac{1}{2}\sum_{j=0}^{n-1}\binom{n+1}{j}\left\{-\sum_{l=j}^{n-1}E_{l-j}+2\right\}\frac{j!}{(j-k)!}\sum_{\substack{0\leq m\leq j-k \\ m\equiv 0 (\operatorname{mod} 2)}}\binom{j-k}{m}E_{j-k-m}\frac{m!}{2^{m+k}k!\left(\frac{m}{2}\right)!} \\
&\quad +(n+1)(-1)^{n+k}\sum_{\substack{0\leq s\leq n-k \\ s\equiv 0 (\operatorname{mod} 2)}}\binom{n}{k}\binom{n-k}{s}E_{n-k-s}\frac{s!}{2^{s+k}\left(\frac{s}{2}\right)!}.
\end{aligned}$$

Consequently, we have the following theorem.

Theorem 2.3. *The following identity holds true:*

$$\begin{aligned}
\sum_{k=0}^{n}E_k(x)x^{n-k} &= \sum_{k=0}^{n}\frac{H_k(x)}{2}\sum_{j=0}^{n-1}\binom{n+1}{j}\left\{-\sum_{l=j}^{n-1}E_{l-j}+2\right\}\frac{j!}{(j-k)!} \\
&\quad \times\sum_{\substack{0\leq m\leq j-k \\ m\equiv 0 (\operatorname{mod} 2)}}\binom{j-k}{m}E_{j-k-m}\frac{m!}{2^{m+k}k!\left(\frac{m}{2}\right)!} \\
&\quad +(n+1)\sum_{k=0}^{n}(-1)^{n+k}\sum_{\substack{0\leq s\leq n-k \\ s\equiv 0 (\operatorname{mod} 2)}}\binom{n}{k}\binom{n-k}{s}H_k(x)E_{n-k-s}\frac{s!}{2^{s+k}\left(\frac{s}{2}\right)!}.
\end{aligned}$$

References

[1] A. Bayad, T. Kim, Identities involving values of Bernstein q-Bernoulli and q-Euler polynomials, Russ. J. Math. Phys. 18 (2011), no. 2, 133-143.

[2] T. Kim J. Choi, Y. H. Kim, C. S. Ryoo, On q-Hermite polynomials, Proc. Jangjeon Math. Soc., 14 (2011), No. 2, 215-221.

[3] T. Kim, Some identities on the q-Euler polynomials of higher order and q-Stiriling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, Russ. J. Math. Phys., 16(2009), no. 4, 484-491.

[4] T. Kim, A note on q-Bernstein polynomials, Russ. J. Math. Phys., 18 (2011), 73-82.

[5] T. Kim, The modified q-Euler numbers and polynomials, Adv. Stud. Contemp. Math. 16 (2008), 161–170.

[6] T. Kim, q-Volkenborn integration, Russ. J. Math. Phys. 9 (2002), 288–299.

[7] T. Kim, On p-adic interpolating function for q-Euler numbers and its derivatives, J. Math. Anal. Appl. 339 (2008), 598–608.

[8] T. Kim, q-generalized Euler numbers and polynomials, Russ. J. Math. Phys. 13 (2006), no. 3, 293-298.

[9] T. Kim, The symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials, J. Difference Equ. Appl. 14 (2008), 1267-1277.

[10] D. S. Kim, D. V. Dolgy, T. Kim and S. H. Rim, Some Formulae for the product of Two Bernoulli and Euler Polynomials, Abstr. Appl. Anal., 2012(2012), Art. ID 784307, 14 pages(Article in Press).

[11] D. S. Kim, T. Kim, S. H. Rim and S. H. Lee., Hermite polynomials and their applications associated with Bernoulli and Euler numbers, Discrete Dynamics in Nature and Society, http://www.hindawi.com/journals/ddns/aip/974632/ (Article in Press).

[12] H. Jolany and H. Sharifi, Some results for Apostol-Genocchi Polynomials of higher order, In press in Bulletin of the Malaysian Mathematical Sciences Society vol 36, no.2, 2013.

[13] M. Acikgoz, Y. Simsek, On multiple interpolation function of the Nörlund-type q-Euler polynomials, Abst. Appl. Anal. 2009 (2009), Article ID 382574, 14 pages.

[14] S. H. Rim and J. Jeong, A note on the Modified q-Euler Numbers and polynomials with weight α, International Mathematical Forum, Vol. 6, 2011, no. 65, 3245-3250.

[15] C. S. Ryoo, A note on the weighted q-Euler numbers and polynomials, Adv. Stud. Contemp. Math. 21 (2011), page 47-54

[16] S. Araci, M. Acikgoz, K. H. Park and H. Jolany, On the unification of two families of multiple twisted type polynomials by using p-adic q-integral on $\mathbb{Z}_p$ at $q = -1$, accepted in Bulletin of the Malaysian Mathematical Sciences and Society.

[17] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p-adic q-integral representation on $\mathbb{Z}_p$ associated with weighted q-Bernstein and q-Genocchi polynomials, Abstract and Applied Analysis, Volume 2011, Article ID 649248, 10 pages.

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `mtsrkn@hotmail.com`

Department of Applied Mathematics, Pukyong National University, Busan 608-737, Republic of KOREA
E-mail address: `seo2011@pknu.ac.kr (Corresponding author)`

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `acikgoz@gantep.edu.tr`